




e-ISSN: 2456-6632

This content is available online at AESA

Archives of Agriculture and Environmental Science

Journal homepage: www.aesacademy.org



## ORIGINAL RESEARCH ARTICLE

## Transcriptional activator gene based phylogenetic analysis of dolichos yellow mosaic virus infecting lablab bean (*Dolichos lablab* L.)

Sonam Arya<sup>1</sup>, Rajat Chaudhary<sup>1</sup>, Vaishali<sup>1</sup> and S.K. Awasthi<sup>2</sup>

<sup>1</sup>Department of Biotechnology, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut-250110 (Uttar Pradesh), INDIA

<sup>2</sup>Department of Life Sciences, Chhatrapati Shahu Ji Maharaj University, Kanpur-208024 (Uttar Pradesh), INDIA

\*Corresponding author's E-mail: arya.sonam22@rediffmail.com

## ARTICLE HISTORY

Received: 01 May 2017  
Revised received: 22 May 2017  
Accepted: 26 May 2017

## Keywords

*Dolichos lablab*  
Dolichos yellow mosaic virus  
Phylogenetic analysis  
Transcriptional activator gene

## ABSTRACT

Lablab bean (*Dolichos lablab* L.) is one the important crop, cultivated as vegetable, pulses as it is rich in protein. It is affected from viral disease i.e. dolichos yellow mosaic virus. The causative agent is begomovirus belongs to geminivirus family. Begomovirus contain bipartite genome having two type of DNA-A and DNA-B. DNA- A helps in replication and DNA-B helps in movement. These DNA have six different type of gene coat protein gene, transcriptional activator gene, replication associated gene, replication enhancer gene, pre coat protein involved in different function associated with it. The present investigation was carried out to investigate the transcriptional activator gene based phylogenetic analysis of Dolichos yellow mosaic virus infecting *D. lablab*. This study is based on the transcriptional activator gene which is used in transactivation of genes, contains three conserved domains: a basic domain at N-Terminus, a central DNA binding domain and activator domain. The genome databases of dolichos yellow mosaic virus were taken from NCBI site total six genome was available and were used with Clustal W and CLC BIO were the bioinformatic tools for determining sequence homology among genome present in different geographical location. The absence of functional specificity suggests that all begomovirus contains a common element interacts with cellular proteins of other viruses reveals the phylogenetic analysis with the other species *Dolichos* in the different geographical location.

©2017 Agriculture and Environmental Science Academy

**Citation of this article:** Arya Sonam, Chaudhary Rajat, Vaishali and Awasthi, S.K. (2017). Transcriptional activator gene based phylogenetic analysis of dolichos yellow mosaic virus infecting lablab bean (*Dolichos lablab* L.). *Archives of Agriculture and Environmental Science*, 2(2): 98-104.

## INTRODUCTION

The lablab bean (*Dolichos lablab* L.) is most ancient cultivated crop plants. It is a multipurpose crop grown for pulse, vegetable and forage, widely cultivated due to it is having high nutritive value and having major sources of protein. It has ability of nitrogen fixation which helps in improving soil fertility, and drought resistance (Capoor and Varma, 1948, 1950; Maruthi *et al.*, 2006). In India, the lablab bean is grown in the peninsular region and cultivated to large areas of Karnataka and adjoining districts of Tamil Nadu, Andhra Pradesh and Maharashtra, Uttar Pradesh (Singh *et al.*, 2006, 2012). The lablab bean is extensively cultivated in the various climatic zones for example arid, semi-arid, sub-tropical and humid regions where temperatures vary between 22°C-35°C, low lands and uplands and many types of soils and the pH varying from 4.4

to 7.8. The dried seeds of lablab bean should be boiled in two changes of water earlier eating since they have toxins substance as Cyanogenic glucosides. It improves the soil condition and is relatively drought tolerant; it is also a good cover crop (Raj *et al.*, 1988; Singh *et al.*, 2006, 2012).

In dolichos disease named dolichos yellow mosaic disease and infected by virus named dolichos yellow mosaic virus (DoYMV) and was shown transmitted by the whitefly *Bemisia tabaci* (Capoor and Varma, 1950; Vanitharani *et al.*, 2004, 2005; Yadava *et al.*, 2010). It having symptoms like faint chlorotic specks on leaf lamina, which later develop into bright yellow mosaic patches with Small Island of green tissue. The leaves are seldom deformed, but yields are reduced significantly (Capoor and Varma, 1950; Bisaro, 1996; Zrachya *et al.*, 2006). The viruses

having bipartite genomes require these components differentiated into two diverse particles; consequently more than one virus particle is needed to infect a cell. The total size of the dolichos yellow mosaic virus is 2760 bp. (Maramorosch and Muniyappa, 1981; Muniyappa Muniyappa *et al.*, 2003; Stanley *et al.*, 2005). Its genome is divided into the two parts DNA-A and DNA -B and intergenic region or common region which is common in both the DNA. All the functions related for virus replication, control of gene expression and encapsidation are in coded in DNA-A and gene involved in intra and intercellular movement are encoded in DNA-B. DNA-A contains six different genes named AC1, AC2, AC3, AC4, AV1, AV2 and each gene have different function. AC2 interacts with itself and localizes to the nucleus and self interaction correlates with nuclear localization and efficient activation of transcription (Hayes and Buck, 1989; Yang *et al.*, 2007; Prajapat *et al.*, 2010). Its function as a transcriptional activator, three conserved domains has been recognized in this protein: a basic domain with a nuclear localization signal (NLS) at the N terminus, a central DNA-binding domain with a non classical Zn-finger motif, and an acidic activator domain at C-terminus (Hartitz *et al.*, 1999; Mansoor *et al.*, 2003; Dhakar *et al.*, 2010). A comparison of steady state transcript level and the transcript level determined by nuclear run on assay showed that activation of AV1 and BV1 gene expression by the AC2 protein primarily occurs at the transcriptional level. Mutational analysis of AC2 in begomoviruses showed that the mutation of this ORF prevented systemic movement of the virus, produced no capsid protein and accumulated reduced amount of ssDNA in the transient assay (Hanley-Bowdoin *et al.*, 1999; Harrison and Robinson, 1999; Markham *et al.*, 1994; Qazi *et al.*, 2007). Therefore, AC2 is functionally interchangeable among begomoviruses and the absence of active specificity suggests that either all begomovirus late promoters have a collective sequence element documented by AC2 or AC2 interacts with cellular proteins common to all begomovirus plant hosts to effect transcriptional activation. AC4 is highly variable among begomoviruses. AC4 of ACMV enhanced pathogenicity by increasing the levels of DNA accumulation of viruses and was also found to have anti-PTGS activity (Varma *et al.*, 1993; Vanitharani *et al.*, 2004; Yang *et al.*, 2007). They infect a wide range of plant species and are responsible for Geminiviruses are a major constraint to agricultural productivity in all tropical and sub-tropical regions of the world. The molecular phylogenetic analysis is the scrutiny of hereditary molecular alterations, mainly in DNA sequences, to gain evidence on an organism's evolutionary relationships (Yaraguntaiah and Govindu, 1964; Briddon *et al.*, 1990, 2001; Varma and Malathi, 2003). The result of a molecular phylogenetic analysis is expressed in a phylogenetic tree (Dry *et al.*, 1997; Swanson *et al.*, 1992; Czosnek *et al.*, 2001). The tree was constructed by use of the various type of software. The CLUSTAL-W, CLC-BIO and BLAST were used to study the phylogenetic analysis. The multiple sequence alignment and phylogenetic relationship was done by the help of CLUSTAL-W and CLC-BIO and BLAST is used

for the global sequence alignment helps in determining the maximum similarity within the other viruses (Rojas *et al.*, 2001; Rekha *et al.*, 2005; Saeed *et al.*, 2008; Govindappa *et al.*, 2011). Keeping in view, the present study was aimed to assess the transcriptional activator gene based phylogenetic analysis of Dolichos yellow mosaic virus infecting Lablab bean (*Dolichos lablab* L.).

## MATERIALS AND METHODS

**Description of study materials:** There are different type's databases available on internet. The NCBI site was used for retrieving the genome of the dolichos yellow mosaic virus (NCBI, 2016). A total of six isolates of the gene AC2 and AC4 were identified for the diversity analysis (Table 1). **Phylogenetic analysis:** Sequences were saved in the FASTA format in a notepad file. Software CLUSTAL-W and CLC-BIO was used for the phlogenetic analysis and making dendrogram. The neucleotide BLAST was done to determine the maximum similarity with the other species. The nucleotide sequences of the AC2 gene in the FASTA format were as follows:

```
>gi|41057586:c1631-1203 Dolichos yellow mosaic virus,
complete genome
ATGCGGTCTTCGTCACCCTCGAACAGCCATT-
GTTCTTACCGAGCATCAAGGCACAGCATCGACAG
GCGAAGAAGCGCAAACGATTCGGCGTAAGCGA
ATCGATCTGACCTGTGGCTGTTTCGTATTACCTAAA
TATCAACTGCCGGAATGACGGATTCACGCACAGG
GGAATCCATCACTGCAACTCAACTACAGAGTGGC
GTGTTTATCTGGGAGCTAAAGAATCCCCTGTCTTT
CAAATCCTGGAGCATCACGAGGGGTTCTGTTC
GATCCGAACCAACACAGAACCAAGATCCGCATAA
TGTTCAACCACGGGTTGAAGAAAGCGTTGGGGAT
ACACAAGGCATTCTTGGATTTTGTGATTTACCATC
GATTGACTCCGCAATCTGGGCGGATCTTGAATGTA
TTCAGGAACTTCCTTTTAA
>gb|AY309241.1|:c1631-1203 Dolichos yellow mosaic
virus segment DNA-A, complete sequence
ATGCGGTCTTCGTCACCCTCGAACAGCCATT-
GTTCTTACCGAGCATCAAGGCACAGCATCGGACAG
GCGAAGAAGCGCAAACGATTCGGCGTAAGCGA
ATCGATCTGACCTGTGGCTGTTTCGTATTACCTGAA
TATCAACTGCCGAAATGACGGATTCACGCACAGG
GGAATCCATCACTGCAGCTCAACTACAGAGTGGC
GTGTTTATTTGGGAGCTAAAGAATCCCCTGTCTTT
CAAATCCTGGAGCATCACGAGGGGTTCTGTTC
GATCCGAACCAACACAGAACCAAGATCCGCATAA
TGTTCAACCACGGGTTGAAGAAAGCGTTGGGGAT
ACACAAGGCATTCTTGGATTTTGTGATTTACCATC
GATTGACTCCGCAATCTGGGCGGATCTTGAATGTA
TTCAGGAACTTCCTTCTTAG
>gb|AY271891.1|:c1631-1203 Hyacinth bean yellow
mosaic Bangladesh virus DNA-A, complete sequence
ATGCGGTCTTCGTCACCCTCGAACAGCCATT-
GTTCTTACCGAGCATCAAGGCACAGCATCGACA
GGCGAAGAAGCGCAAACGATTCGGCGTAAGCG
AATCGATCTGACCTGTGGCTGTTTCGTATTACCTAA
ATATCAACTGCCGGAATGACGGATTCACGCACAG
GGGAATCCATCACTGCAACTCAACTACAGAGTGG
```

CGTGTTTATCTGGGAGCTAAAGAATCCCCTGTCTT  
TCAAAATCCTGGAGCATCACGAGGGGTTCTGT  
CGATCCGAACCAACACAGAACCAAGATCCGCATA  
ATGTTCAACCACGGGTTGAAGAAAGCGTTGGGGA  
TACACAAGGCATTCTTGGATTTTGTGATTTACCAT  
CGATTGACTCCGCAATCTGGGCGGATCTTGAATGT  
ATTCAGGAACCTCCTTTTTAA

>gi|116077869:c1631-1203 Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolate Bangalore-2  
ATGCGGTCTTCGTCACCCTCGAACAGCCATT-  
GTTCTCTACCGAGCATCAAGGCACAGCATCGACA  
GGCGAAGAAGCGCAAACGATTTCGGCGTAAGCG  
AATCGATCTGACCTGTGGCTGTTTCGTATTACCTAA  
ATATCAACTGCCGGAATGACGGATTCACGCACAG  
GGGAATCCATCACTGCAGCTCAACTACAGAGTGG  
CGTGTTTATCTGGGAGCTAAAGAATCCCCTGTCTT  
TCAAAATCCTGGAGCATCACGAGGGGTTCTGT  
CGATCCGAACCAACACAGAACCAAGATCCGCATA  
ATGTTCAACCACGGGTTGAAGAAAGCGTTGGGGA  
TACACAAGGCATTCTTGGACTTAGTAGTTTACCAT  
CGATTGACTCCGCAATCTGGGCGGATCTTGAATGT  
ATTCAGGAACCTCCTTTTTAG

>gi|116077862:c1631-1203 Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolate Bangalore-1  
ATGCGGTCTTCGTCACCCTCGAACAGCCATT-  
GTTCTCTACCGAGCATCAAGGCACAGCATCGACA  
GGCGAAGAAGCGCAAACGATTTCGGCGTAAGCG  
AATCGATCTGACCTGTGGCTGTTTCGTATTACCTAA  
ATATCAACTGCCGGAATGACGGATTCACGCACAG  
GGGAATCCATCACTGCAGCTCAACTACAGAGTGG  
CGTGTTTATCTGGGAGCTAAAGAATCCCCTGTCTT  
TCAAAATCCTGGAGCATCACGAGGGGTTCTGT  
CGATCCGAACCAACACAGAACCAAGATCCGCATA  
ATGTTCAACCACGGGTTGAAGAAAGCGTTGGGGA  
TACACAAGGCATTCTTGGACTTAGTAGTTTACCAT  
CGATTGACTCCGCAATCTGGGCGGATCTTGAATGT  
ATTCAGGAACCTCCTTTTTAG

>gi|90185977:c1631-1203 Dolichos yellow mosaic virus complete DNA-A segment, strain Mysore  
AATCGGTCTTCGTCACCCTCGAACAGCCATT-  
GTTCTCTACCGAGCATCAAGGCACAGCATCGACA  
GGCGAAGAAGCGCAAACGATTTCGGCGTAAGCG  
AATCGATCTGACCTGTGGCTGTTTCGTATTACCTAA  
ACATCAACTGCCGGAATGACGGATTCACGCACAG  
GGGAATCCATCACTGCAGCTCAACTACAGAGTGG  
CGTGTTTATCTGGGAGCTAAAGAATCCCCTGTCTT  
TCAAAATCCTGGAACATCACGAGGGGTTCTGT  
CGATCCGAACCAACACAGAACCAAGATCCGCATA  
ATGTTCAACCACGGGTTGAAGAAAGCGTTGGGGA  
TACACAAGGCATTCTTGGACTTTGTGATTTACCAT  
CGATTGACTCCGCAATCTGGGCGGATCTTGAATGT  
ATTCAGGAACCTCCTTTTTAA

**Nucleotide sequence of AC4 gene in the FASTA format:** >gi|41057586:c2480-2235 Dolichos yellow mosaic virus, complete genome

ATGGGACTATGCATCTCCATGCCCTCGTCCAG-  
TTCAAGGGTAAGGCCAGTTCCGAAATGCAAGAC  
ATTTGCACCTTACCCATCCTCATAGCACACAGGTT  
TTCCATGGAAATGTCCAGGGAGCAAAGAGCTCAT

CTGATGTCAAATCCTACATCACCAAGGACGGTGA  
TTACGTGCGACTGGGGAACATTTTCAGATCGACGGA  
CGATCTGCTAGAGGAGGTCGTCAGACAGCTGACG  
ATGCTGTAG

>gb|AY309241.1|:c2480-2235 Dolichos yellow mosaic virus segment DNA-A, complete sequence

ATGGGA-  
TATGCATCTCCATGCCCTCGTCCATTCAAGGGTAA  
GGCCCAGTTCCGAAATGCAAGACATTTTCGACCTT  
ACCCATCCTCATAGCACACAGGTTTTCCATGGAAA  
TGTCAGGGAGCAAAGAGCTCATCTGATGTCAA  
TCCTACATCACCAAGAACGGTGATTACATCGACT  
GGGGAACATTTTCAGATCGACGGACGATCTGCTAG  
AGGAGGTCGTCAGACAGCTGACGATGCTGTAG

>gb|AY271891.1|:c2480-2235 Hyacinth bean yellow mosaic Bangladesh virus DNA-A, complete sequence

ATGGGA-  
TATGCATCTCCATGCCCTCGTCCATTCAAGGGTAA  
GGCCCAGTTCCGAAATGCAAGACATTTTCGACCTT  
ACCCATCCTCATAGCACACAGGTTTTCCATGGAA  
ATGTCCAGGGAGCAAAGAGCTCATCTGATGTCAA  
ATCCTACATCACCAAGGACGGTGATTACGTGCGAC  
TGGGGAACATTTTCAGATCGACGGACGATCTGCTA  
GAGGAGGTCGTCAGACAGCTGACGATGCTGTAG

>gi|116077869:c2480-2235 Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolates Bangalore-2

ATGGGACTATGCATCTCCATGCCCTCGTCCAG-  
TTCAAGGGTAAGGCCAGTTCCGAAATGCCAGAC  
ATTTGCACCTTACCCATCCTCATAGCACACAGATT  
TTCCATGGAAATGTCCAGGGAGCAAAGAGCTCAT  
CTGATGTCAAATCCTACATCACCAAGGACGGTGA  
CTACGTGCGACTGGGGAACATTTTCAGATCGACGGA  
CGACCTGCTAGAGGAGGTCGGCAGACAGCTGACG  
ATGCTGTAG

>gi|116077862:c2480-2235 Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolate Bangalore-1

ATGGGACTATGCATCTCCATGCCCTCGTCCAG-  
TTCAAGGGTAAGGCCAGTTCCGAAATGCCAGAC  
ATTTGCACCTTACCCATCCTCATAGCACACAGATT  
TTCCATGGAAATGTCCAGGGAGCAAAGAGCTCAT  
CTGATGTCAAATCCTACATCACCAAGGACGGTGA  
CTACGTGCGACTGGGGAACATTTTCAGATCGACGGA  
CGACCTGCTAGAGGAGGTCGGCAGACAGCTGACG  
ATGCTGTAG

>gi|90185977:c2480-2235 Dolichos yellow mosaic virus complete DNA-A segment, strain Mysore

GATGGGA-  
TATGCATCTCCATGCCCTCGTCCATCAAGGGTAA  
GGCCCAGTTCCGAAATGCCAGACATTTTCGACCTTAC  
CCATCCTCATAGCACACAGATTTTCCATGGAAATG  
TCCAGGGAGCAAAGAGCTCATCTGATGTCAAATC  
CTACATCACCAAGGACGGTACTACGTGCGACTGG  
GGAACATTTTCAGATCGACGGACGATCTGCTAGAG  
GAGGTCGTCAGACAGCTGACGATGCTGTA

## RESULTS AND DISCUSSION

The yellow mosaic diseases are challenging problems in the tropics and subtropics. The increase in *Bemisia tabaci*

**Table 1.** Isolates used for phylogenetic study.

S.N.	Accession No.	Gi.no	Description	Size of the gene	Place of isolate
1	NC-005338.1	41057586	Dolichos yellow mosaic virus isolate DYMV complete genome	AC2-1203-1631 AC4-2235-2480	South and South east Asia
2	AY309241.1	32187104	Dolichos yellow mosaic virus complete DNA-A segment	AC2-1203-1631 AC4-2235-2480	Mysore
3	AM157413.1	116077869	Dolichos yellow mosaic virus DNA-A, complete sequence.	AC2-1203-1631 AC4-2235-2480	Banglore-2
4	AM157412.1	116077862	Dolichos yellow mosaic virus segment DNA-A, complete sequence	AC2-1203-1631 AC4-2235-2480	India
5	AJ968370.1	90185977	Dolichos yellow mosaic virus DNA A, complete genome	AC2-1203-1631 AC4-2235-2480	Mysore
6	AY271891.1	33355890	Hyacinth bean yellow mosaic virus segment DNA-A, complete sequence	AC2-1203-1631 AC4-2235-2480	Bangladesh

**Table 2.** BLAST Score result showing maximum similarity with other viruses.

Accession	Description	Max score	Total score	Query coverage	E value	Max ident	Links
<a href="#">AY271891.1</a>	Hyacinth bean yellow mosaic Bangladesh virus DNA-A, complete sequence	<a href="#">5099</a>	5099	100%	0.0	100%	<a href="#">G</a>
<a href="#">AJ968370.1</a>	Dolichos yellow mosaic virus complete DNA-A segment, strain Mysore	<a href="#">4505</a>	4505	100%	0.0	96%	
<a href="#">AM157413.1</a>	Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolate Bangalore-2	<a href="#">4447</a>	4447	100%	0.0	96%	
<a href="#">AM157412.1</a>	Dolichos yellow mosaic virus DNA-A segment, complete sequence, isolate Bangalore-1	<a href="#">4447</a>	4447	100%	0.0	96%	
<a href="#">AY309241.1</a>	Dolichos yellow mosaic virus segment DNA-A, complete sequence	<a href="#">4385</a>	4385	100%	0.0	95%	<a href="#">G</a>
<a href="#">GU591170.1</a>	Dolichos yellow mosaic virus isolate DYMV-CpKn coat protein (AV1) gene, complete cds	<a href="#">1242</a>	1242	28%	0.0	96%	
<a href="#">JN368437.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Rembang segment DNA-A, complete sequence	<a href="#">407</a>	659	47%	3e-109	80%	
<a href="#">JN368432.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Bogor segment DNA-A, complete sequence	<a href="#">407</a>	652	48%	3e-109	80%	
<a href="#">JN368433.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Purwakarta 1 segment DNA-A, complete sequence	<a href="#">401</a>	652	48%	1e-107	79%	
<a href="#">FN543425.1</a>	Velvet bean severe mosaic virus DNA A, complete genome	<a href="#">401</a>	401	23%	1e-107	78%	<a href="#">G</a>
<a href="#">JN368439.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Brebes 4 segment DNA-A, complete sequence	<a href="#">396</a>	646	48%	6e-106	79%	
<a href="#">JN368434.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Purwakarta 2 segment DNA-A, complete sequence	<a href="#">396</a>	646	48%	6e-106	79%	
<a href="#">JN368436.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Brebes 2 segment DNA-A, complete sequence	<a href="#">390</a>	646	48%	3e-104	79%	
<a href="#">JN368435.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Brebes 1 segment DNA-A, complete sequence	<a href="#">390</a>	641	48%	3e-104	79%	
<a href="#">JN368438.1</a>	Mungbean yellow mosaic India virus Indonesia isolate Brebes 3 segment DNA-A, complete sequence	<a href="#">385</a>	646	48%	1e-102	79%	
<a href="#">AY618902.1</a>	Cowpea golden mosaic virus segment DNA-A, complete sequence	<a href="#">385</a>	635	48%	1e-102	79%	
<a href="#">AY937199.1</a>	Mungbean yellow mosaic India virus clone MB 0.9 segment DNA A, partial sequence	<a href="#">385</a>	385	20%	1e-102	79%	
<a href="#">AY937195.1</a>	Mungbean yellow mosaic India virus clone MBK-A25 segment DNA A, complete sequence	<a href="#">379</a>	635	48%	6e-101	79%	
<a href="#">JN181006.1</a>	Mungbean yellow mosaic India virus isolate Vizianagaram pre-coat protein (AV2) gene, partial cds; and coat protein (AV1) gene, complete cds	<a href="#">374</a>	374	23%	3e-99	77%	
<a href="#">FJ685621.1</a>	Tomato leaf curl Nigeria virus-[Nigeria:2006], complete sequence	<a href="#">372</a>	372	22%	9e-99	78%	<a href="#">G</a>
<a href="#">JQ398669.1</a>	Mungbean yellow mosaic virus [Urdbean:New Delhi:2011] clone MF2 segment A, complete sequence	<a href="#">370</a>	628	47%	3e-98	78%	
<a href="#">HE616781.1</a>	African cassava mosaic Burkina Faso virus DNA-A complete genome, isolate BF:Oua:BF128C:08	<a href="#">368</a>	368	22%	1e-97	78%	
<a href="#">HE616780.1</a>	African cassava mosaic Burkina Faso virus DNA-A complete genome, isolate BF:Oua:BF128B:08	<a href="#">368</a>	368	22%	1e-97	78%	
<a href="#">HE616777.1</a>	African cassava mosaic Burkina Faso virus DNA-A complete genome, isolate BF:Oua:BF127:08	<a href="#">368</a>	368	22%	1e-97	78%	
<a href="#">HE616779.1</a>	African cassava mosaic Burkina Faso virus DNA-A complete genome, isolate BF:Oua:BF128A:08	<a href="#">363</a>	363	22%	6e-96	78%	
<a href="#">FM208839.1</a>	Mungbean yellow mosaic India virus, segment DNA-A, complete sequence, clone A14	<a href="#">353</a>	624	47%	3e-93	78%	
<a href="#">AY049772.1</a>	Mungbean yellow mosaic India virus-[Soybean] segment DNA A, complete sequence	<a href="#">353</a>	353	20%	3e-93	78%	
<a href="#">GQ387509.1</a>	Mungbean yellow mosaic India virus isolate MYMIV-Ub04 coat protein (AV1) gene, complete cds	<a href="#">351</a>	351	20%	1e-92	78%	
<a href="#">AM992618.1</a>	Mungbean yellow mosaic India virus, segment DNA-A, complete genome	<a href="#">351</a>	351	23%	1e-92	77%	




Figure 1. Phylogenetic tree of AC2 gene created by CLC-BIO.




Figure 2. Phylogenetic tree of AC4 gene created by CLC-BIO.




Figure 3. Multiple sequence alignment result of AC2 gene.

populations during the cropping season has also led to geminivirus problems in other countries evolution of new variants of the viruses depends upon the variability among viruses I, it arises through mutations, development of recombinants and pseudo recombinants. In the present study, we examine the influence of variability in Gemini viruses in relation to new disease problems. Variability in Gemini viruses is seen especially between isolates from geographically different regions. This has been shown for MSV, where emerging *Gemini viruses* 6 isolates genomic recombination in *Gemini viruses* may occur not only between variants of the same virus but also between virus species (Tables 1, 2). In the above case, six different evolutionary pathways are possible using an out group, each depicted by a different rooted tree. The sequence of one clone was determined to be 2760 nucleotides in length (accession number AJ968370) and to be most similar to the sequences of DoYMV-[Bangladesh] (AY271891) and a further clone of DoYMV originating from India (AY309241) at 95.2% and 93.4% nucleotide sequence identities, respectively. The three DoYMV isolates examined in this study showed the highest levels of sequence identity (92.5–95.3%) with the previously described DoYMV




Figure 4. Multiple sequence alignment result of AC4 gene.

isolates from North India and Bangladesh. DoYMV-[Ban1] and DoYMV-[Ban2] originated from two separate fields in Bangalore and were almost identical (99.1%) (Table 2). Iiyas *et al.* (2010) reported genetic diversity and phylogeography of begomoviruses infecting legumes in Pakistan. Moreover Maruthi *et al.* (2006) also reported genetic similarities in the different begomovirus causes Indian dolichos yellow mosaic disease. In another study Maruthi *et al.* (2006) also concluded that dolichos yellow mosaic virus belongs to a distinct lineage of Old World begomoviruses and it showed variations in its biological and molecular properties. During the present study, the sequences of the genomes (DNA-A components) of the two DoYMV isolates originating from Bangalore (DoYMV-[Ban1] and DoYMV-[Ban2]) were both 2762 nucleotides in length whereas the length of DoYMV-[Mys] was 2760 nucleotides it has been noted previously that the antigenic and genomic variation in begomoviruses is primarily correlated with geography rather than their host range. Viruses infecting different host plants but originating from the same geographical area are more likely to be closely related than viruses infecting the same host but from different

geographical areas Dolichos yellow mosaic disease was suspected to be caused by a geminivirus after the successful transmission of the pathogen from DYMD-affected plants by *B. tabaci*. The phylogenetic trees and multiple sequence alignment result of the gene AC2 and AC4 showed the geographical similarities (Figures 1-4). The findings are in agreement with Padidam *et al.* (1996) who reported the role of AV2 (precoat) and coat protein in viral replication and movement in tomato leaf curl geminivirus. In an almost similar study Muniyappa *et al.* (1987) also isolated and characterized of Dolichos yellow mosaic disease geminivirus causing yellow mosaic disease of Horsegram (*Macrotyloma uniflorum* (Lam.) Verdc.).

### Conclusions

The present study concluded that variations in Gemini viruses were seen especially between isolates from geographically different regions and this is likely due to the MSV, where *Emerging Gemini viruses 6* Isolates Genomic recombination in Gemini viruses may occur not only between variants of the same virus but also between virus species. The sequence of one clone was determined to be 2760 nucleotides in length (accession number AJ968370) and to be most similar to the sequences of DoYMV-[Bangladesh] (AY271891) and a further clone of DoYMV originating from India (AY309241) at 95.2% and 93.4% nucleotide sequence identities, respective. The three DoYMV isolates examined in this study showed the highest levels of sequence identity (92.5–95.3%) with the previously described DoYMV isolates from North India and Bangladesh. DoYMV-[Ban1] and DoYMV-[Ban2] originated from two separate fields in Bangalore and were almost identical (99.1%). The sequences of the genomes (DNA-A components) of the two DoYMV isolates originating from Bangalore (DoYMV-[Ban1] and DoYMV-[Ban2]) were both 2762 nucleotides in length whereas the length of DoYMV-[Mys] was 2760 nucleotides it has been noted previously that the antigenic and genomic variation in begomoviruses is primarily correlated with geography rather than their host range. Therefore, viruses infecting different host plants but originating from the same geographical area are more likely to be closely related than viruses infecting the same host but from different geographical areas Dolichos yellow mosaic disease was suspected to be caused by a geminivirus.

**Open Access:** This is open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

### REFERENCES

- Bisaro, D.M. (1996). Geminivirus DNA replication. In DNA Replication in eukaryotic Cells, pp. 833-854. Edited by M. L. DePamphilis. Cold Spring Harbor, NY.
- Briddon, R.W., Mansoor, S., Bedford, I.D., Pinner, M.S., Saunders, K., Stanley, J., Zafar, Y., Malik, K. and Markham, P.G. (2001). Identification of DNA components required for induction of cotton leaf curl disease. *Virology*, 285: 234-243.
- Briddon, R.W., Pinner, Mansoor, S., Stanley, J. and Markham, P.G. (1990). Geminivirus coat protein replacement alters insect specificity. *Virology*, 177: 85-94.
- Capoor, S.P. and Varma, P.M. (1948). Yellow mosaic of *Phaseolus lunatus* L. *Current Science*, 17: 152-153.
- Capoor, S.P. and Varma, P.M. (1950). New virus disease of *Dolichos lablab*. *Current Science*, 19: 242-249.
- Czosnek, H., Ghanim, M., Morin, S., Rubinstein, G., Fridman, V. and Zeidan, M. (2001). Whiteflies; vectors, and victims of geminiviruses. *Advances in Virus Research*, 57: 291-322.
- Dhakar, K., Gupta, V.K., Rathore, M.S. and Gaur, R.K. (2010). Virus Resistance and Gene Silencing in Plants Infected with Begomovirus. *Journal of Applied Sciences*, 10: 1787-1791.
- Dry, I., Krake, L.R., Rigden, J.E. and Rezaian, M.A. (1997). A novel subviral agent associated with a geminivirus: the first report of a DNA satellite. *Proceedings of the National Academy of Sciences USA*, 94: 7088-7093.
- Govindappa, M.R. Shankergoud, I., Shankarappa, K. S. and Wickramaarachchi, W.A.R.T. (2011). Molecular Detection And Partial Characterization of Begeomovirus Associated With Leaf Curl Disease Of Sun Flower (*Helianthus annuus*) in Southern India. *Plant Pathology Journal*, 10 (1): 29-35.
- Hanley-Bowdoin, L., Settlege, S.B., Orozoo, B.M., Nagar, S. and Robertson, D. (1999). Geminiviruses: models for plant DNA replication, transcription, and cell cycle regulation. *Critical Reviews in Plant Sciences*, 18: 71-106.
- Harrison, B.D. and Robinson, D.J. (1999). Natural genomic and antigenic variation in whitefly-transmitted geminiviruses. *Annual Review of Phytopathology*, 37: 369-398.
- Hartz, M.D., Sunter, G. and Bisaro, D.M. (1999). The geminivirus transactivator (TrAP) is a single-stranded DNA and zinc-binding phosphoprotein with an acidic activation domain. *Virology*, 263: 1-14.
- Hayes, R.J. and Buck, K.W. (1989). Replication of tomato golden mosaic virus DNA B in transgenic plants expressing open reading frames (ORFs) of DNA A: requirement of ORF AL2 for production of single-stranded DNA. *Nucleic Acids Research*, 17: 10213-10222.
- Iiyas, Muhammad, Qazi, Javaria, Mansoor, Shahid and Briddon, W.R. (2010). Genetic diversity and phylogeography of begomoviruses infecting legumes in Pakistan. *Journal of General Virology*, 91: 2091-2101.
- Maruthi, M.N., Rekha, A.R., Govindappa, M.R., Colvin, J. and Muniyappa, V. (2006). A distinct begomovirus causes Indian dolichos yellow mosaic disease. *Plant Pathology*, 55: 290.
- Mansoor, S., Briddon, R. W., Zafar, Y., and Stanley, J. (2003). Geminivirus disease complexes: an emerging threat. *Trends Plant Science*, 8: 128-134.
- Maramorosch, K. and Muniyappa, V. (1981). Whitefly transmitted plant disease agents in Karnataka, India. In International Workshop on Pathogens Transmitted by Whiteflies, 31 July 1981, Association of Applied Biologists, Oxford, UK.
- Naraini T.K. (1960) Yellow mosaic of mung (*Phaseolus aureus*). *Indian Phytopathology*, 13: 24-29.
- Markham, P.G., Bedford, I.D., Liu, S. and Pinner, M.S. (1994). The transmission of geminiviruses by *Bemisia tabaci*. *Pesticide Science*, 42: 123-128.
- Muniyappa, V., Rajeshwari, R., Bharathan, N., Reddy, D.V.R. and Nolt, B.L. (1987). Isolation and characterization of Dolichos yellow mosaic disease geminivirus causing yellow mosaic disease of Horsegram (*Macrotyloma uniflorum* (Lam.) Verdc.). *Indian Journal of Phytopathology*, 119: 81-87.

- Muniyappa, V., Maruthi, M.N., Babitha, C.R., Colvin, J., Briddon, R.W. and Rangaswamy, K.T. (2003). Characterization of pumpkin yellow vein mosaic virus from India. *Annals of Applied Biology*, 142: 323-331.
- Maruthi, M.N., Manjunatha, B., Rekha, A.R., Govindappa, M.R., Colvin, J., and Muniyappa, V. (2006). Dolichos yellow mosaic virus belongs to a distinct lineage of Old World begomoviruses; its biological and molecular properties. *Annals of Applied Biology*, 1744-1747.
- NCBI (2016). Dolichos yellow mosaic virus. Retrieved from <https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=333968>.
- Padidam, M., Beachy, R. N., and Fauquet, C. M. (1996). The role of AV2 (precoat) and coat protein in viral replication and movement in tomato leaf curl geminivirus. *Virology*, 224: 390-404.
- Prajapat, R., Gaur, R.K., Raizada, R. and Gupta, V.K. (2010). In silico analysis of genetic diversity of begomovirus using homology modeling. *Journal of Biological Sciences*, 217-223.
- Qazi, Javaria, Iiyas, Muhammad, Mansoor, Shahid and Briddon, W.R. (2007). Legume yellow mosaic viruses: genetically isolated begomoviruses. *Molecular Plant Pathology*, 343-348.
- Raj, S.K., Aslam, N., Srivastava, K.M. and Singh, B.P. (1988). Association of Geminivirus-like particles with yellow mosaic disease of *Dolichos lablab*. *Current Science*, 58: 813-814.
- Rekha, A.R., Maruthi, M.N., Muniyappa, V. and Colvin, J. (2005). Occurrence of three genotypic clusters of *Bemisia tabaci* (Gennadius) and the rapid spread of the B-biotype in South India. *Entomologia Experimentalis et Applicata*, 117: 221-233.
- Rojas, M.R., Jiang, H., Salati, R., Xoconostle-Cazares, B., Sudarshana, M.R., Lucas, W.J. and Gilbertson, R.L. (2001). Functional analysis of proteins involved in movement of the monopartite begomovirus, Tomato yellow leaf curl virus. *Virology*, 291 110-125.
- Saeed, M., Mansoor, S., Rezaian, M.A., Briddon, R.W. and Randles, J.W. (2008). Satellite DNA  $\beta$  overrides the pathogenicity phenotype of the C4 gene of Tomato leaf curl virus, but does not compensate for loss of function of the coat protein and V2 genes. *Archives of Virology*, 153: 1367-1372.
- Singh, S.K., Chakraborty, S., Singh, A.K. and Pandey, P.K. (2006). Cloning, restriction mapping and phylogenetic relationship of genomic components of MYMIV from *Lablab purpureus*. *Bioresource Technology*, 97: 1807-1814.
- Singh, P.K., Rai, N., Singh, D.V. and Singh, A.P. (2012). Incidence of Dolichos yellow mosaic virus (DYMV) and yield potential in Indian bean (*Lablab purpureus*) F1'S. *Journal of Agricultural Technology*, 8(4): 1469-1474.
- Swanson, M.M., Varma, A., Muniyappa, V. and Harrison, B.D. (1992). Comparative epitope profiles of the particle proteins of whitefly-transmitted geminiviruses from nine crop legumes in India. *Annals of Applied Biology*, 120: 425-433.
- Stanley, J., Bisaro, D.M., Briddon, R.W., Brown, J.K., Fauquet, C.M., Harrison, B.D., Rybicki, E.P. and Stenger, D.C. (2005). *Geminiviridae* In: Fauquet, C.M., Mayo, M.A., Maniloff, J., Desselberger, U., Ball, L.A. (Eds), VIIIth Report of the International Committee on Taxonomy of Viruses. Virus Taxonomy. Elsevier/Academic Press, London, pp. 1163-1169.
- Varma, A. and Malathi, V.G. (2003). Emerging geminivirus problems. A serious threat to crop production. *Annals of Applied Biology*, 142: 145-164.
- Varma, A., Malathi, V. G., Handa, A., Aiton, M., Harrison, B. D. Varma, J.P., Singh, R.P., Singh, M., Srivastava, M. and Singh, J. (1993). Occurrence of leaf-curl of cotton and okra in Northern India. Abstracts of the 6th International Congress of Plant Pathology, Montreal, Canada, pp. 17: 5-14.
- Vanitharani, R., Chellappan, P., Pita, J.S., Fauquet, C.M. (2004). Differential roles of AC2 and AC4 of cassava geminiviruses in mediating synergism and suppression of posttranscriptional gene silencing. *Journal of Virology*, 78: 9487-9498.
- Vanitharani, R., Chellappan, P. and Fauquet, C. M. (2005). Geminiviruses and RNA silencing. *Trends in Plant Science*, 10: 144-151
- Yadava, P., Suyal, G. and Mukherjee, S.K. (2010). Begomovirus DNA replication and pathogenicity. *Current Science*, 98:3-10.
- Yang, X., Baliji, S., Buchmann, R.C., Wang, H., Lindbo, J.A., Sunter, G., and Bisaro, D.M. (2007). Functional modulation of the geminivirus AL2 transcription factor and silencing suppressor by self-interaction. *Journal of Virology*, 81: 11972-11981.
- Yaraguntaiah, R.C. and Govindu, H.C. (1964). Virus disease of *Dolichos lablab* var. typicum from Mysore. *Current Science*, 33: 721-722.
- Zracha, A., Glick, E., Levy, Y., Arazi, T., Citovsky, V. and Gafni, Y. (2006). Suppressor of RNA silencing encoded by tomato yellow leaf curl virus-Israel. *Virology*, 358: 159-165.